arXiv:1304.0214v3 [math.DS] 24 Apr 2013

Ergodic Properties of m-Free Integers in Number Fields

Francesco Cellarosi[*], Ilya Vinogradov[†]

April 25, 2013

Abstract

Let $K/\mathbf{Q}$ be a degree d extension. Inside the ring of integers $\mathcal{O}_K$ we define the set of m-free integers $\mathcal{F}_m$ and a natural $\mathcal{O}_K$-action on the space of binary $\mathcal{O}_K$-indexed sequences, equipped with a natural $\mathcal{O}_K$-invariant probability measure naturally associated to $\mathcal{F}_m$. We prove that this action is ergodic, has pure point spectrum and is isomorphic to a $\mathbf{Z}^d$-action on a compact abelian group. In particular, it is not weakly mixing and has zero measure-theoretical entropy. This work generalizes [3] where $K = \mathbf{Q}$ and $m = 2$.

Keywords: square-free and m-free integers in number fields, correlation functions, group actions with pure point spectrum, ergodicity, isomorphism of group actions. **MSC**: 37A35, 37A45, 11R04, 11N25, 37C85, 28D15.

Introduction and Notation

Let K be a number field of degree $[K : \mathbf{Q}] = d$, and let $\mathcal{O}_K$ be its ring of integers. While $\mathcal{O}_K$ need not be a principal ideal domain, it is always a Dedekind domain, that is an integral domain whose proper ideals factor (uniquely, up to the order) into a product of prime ideals (see [9, 10]). We will denote ideals in $\mathcal{O}_K$ by $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}, \mathfrak{n}$, and prime ideals by $\mathfrak{p}, \mathfrak{q}$. The sum of two ideals is defined as $\mathfrak{a} + \mathfrak{b} = \{a + b : \ a \in \mathfrak{a},\ b \in \mathfrak{b}\}$, the product as $\mathfrak{a}\mathfrak{b} = \{\sum_{i=1}^{n} a_i b_i,\ a_i \in \mathfrak{a},\ b_i \in \mathfrak{b},\ n \in \mathbf{N}\}$, and these are ideals. We say that $\mathfrak{a}$ *divides* $\mathfrak{b}$ if and only if $\mathfrak{b} = \mathfrak{a}\mathfrak{c}$ for some ideal $\mathfrak{c}$ or, equivalently, if and only if $\mathfrak{a} \supseteq \mathfrak{b}$. For any two ideals $\mathfrak{a}, \mathfrak{b}$ we define their *greatest common divisor* as the smallest ideal containing both $\mathfrak{a}$ and $\mathfrak{b}$, that is $\gcd(\mathfrak{a}, \mathfrak{b}) = \mathfrak{a} + \mathfrak{b}$. The *least common multiple* of $\mathfrak{a}$ and $\mathfrak{b}$ is defined as the largest ideal contained in both $\mathfrak{a}$ and $\mathfrak{b}$,

---

[*] University of Illinois at Urbana-Champaign. IL, U.S.A. `fcellaro@illinois.edu`

[†] University of Bristol. U.K. `ilya.vinogradov@bristol.ac.uk`

that is $\operatorname{lcm}(\mathfrak{a},\mathfrak{b}) = \mathfrak{a}\cap\mathfrak{b}$. Let $m \geqslant 2$. An ideal $\mathfrak{a}$ is *m-free* if $\mathfrak{p}^m \not\supseteq \mathfrak{a}$ for every prime ideal $\mathfrak{p}$. For a nonzero ideal $\mathfrak{a}$, the *Möbius function* is defined as

$$\mu(\mathfrak{a}) = \begin{cases} 1 & \text{if } \mathfrak{a} = \mathcal{O}_K; \\ 0 & \text{if } \mathfrak{a} \text{ is not square-free}; \\ (-1)^k & \text{if } \mathfrak{a} \text{ is the product of } k \text{ distinct prime ideals.} \end{cases}$$

In particular $\mathfrak{a}$ is square-free if and only if $\mu^2(\mathfrak{a}) = 1$. Let μ_m be the indicator of m-free ideals. In particular $\mu_2 = \mu^2$. In general, we have

$$\mu_m(\mathfrak{a}) = \sum_{\mathfrak{b}^m \supseteq \mathfrak{a}} \mu(\mathfrak{b}). \tag{1}$$

We will say that two integers $a, b \in \mathcal{O}_K$ are *congruent modulo the ideal* $\mathfrak{a}$ (denoted by $a \equiv b \mod \mathfrak{a}$) iff $a + \mathfrak{a} = b + \mathfrak{a}$ in the finite ring $\mathcal{O}_K/\mathfrak{a}$. The *(algebraic) norm*[1] of a nonzero ideal $\mathfrak{a}$ is defined as $N(\mathfrak{a}) = \#\mathcal{O}_K/\mathfrak{a} = [\mathcal{O}_K : \mathfrak{a}]$, where $\#\cdot$ denotes the cardinality of a finite set. The *Dedekind zeta function* is given, for $\Re s > 1$, by

$$\zeta_K(s) = \sum_{\mathfrak{a}} N(\mathfrak{a})^{-s} = \prod_{\mathfrak{p}} \left(1 - N(\mathfrak{p})^{-s}\right)^{-1},$$

where the sum ranges over all nonzero ideals of $\mathcal{O}_K$ and the product over the prime ones.

If $a \in \mathcal{O}_K$, we set $\mu(a) = \mu((a))$ and $N(a) = N((a))$, where (a) denotes the ideal generated by a. Moreover, $a \in \mathcal{O}_K$ is said to be *m-free* if (a) is m-free. The set of m-free integers is denoted by $\mathcal{F}_m$.

Henceforth, let us fix d generators for $\mathcal{O}_K$, i.e. elements $e_1, \ldots, e_d \in \mathcal{O}_K$ such that $\mathbf{Z}[e_1, \ldots, e_d] = \mathcal{O}_K$, and an isomorphism $\iota : \mathbf{Z}^d \to \mathcal{O}_K$, $(\alpha_1, \ldots, \alpha_d) \mapsto \alpha_1 e_1 + \ldots + \alpha_d e_d$. None of our results will depend on this choice except for implied constants in error terms. For an element $a \in \mathcal{O}_K$ take $\operatorname{Size} a$ to be defined as the distance to 0 in the resulting Cayley graph (other notions of size will also work[2]). Let $B_x = \{a \in \mathcal{O}_K : \operatorname{Size} a \leqslant x\}$ denote the ball or radius x in $\mathcal{O}_K$, with respect to Size. Suppose $\mathfrak{a} = \langle \sum_j k_{ij} e_j \rangle_{i=1}^d = \langle v_i \rangle_{i=1}^d$. For any choice of generators we call the set $\Delta = \{\sum_j \varepsilon_{ij} v_j : \varepsilon = 0 \text{ or } 1\}$ a *cell* of $\mathfrak{a}$. The *diameter* of an ideal $\mathfrak{a}$, written $\operatorname{diam} \mathfrak{a}$, is defined by

$$\min_{\Delta \text{ is a cell of } \mathfrak{a}} \left(\max_{a \in \Delta} \operatorname{Size} a \right).$$

1 Statement of results

Fix $m \geqslant 2$. The goal of this paper is to describe a $\mathcal{O}_K$-action 'naturally' associated to $\mathcal{F}_m$ and study its statistical and ergodic properties.

---

[1] There is also a notion of *norm* for elements of K. Let $a \in K$ and let $m_a : K \to K$ be the $\mathbf{Q}$-linear map $m_a(b) = ab$. The norm $N_{K/\mathbf{Q}}(a)$ is defined as the determinant of m_a. If $K/\mathbf{Q}$ is Galois, then $N_{K/\mathbf{Q}}$ equals the product of the conjugates of a. When $\mathcal{O}_K$ is a PID, and $\mathfrak{a} = (a)$, then $N(\mathfrak{a}) = N_{K/\mathbf{Q}}(a)$.

[2] For example, any (vector space) norm on $\mathbf{R}^d$ induces one on $\mathbf{Z}^d$ by restriction and hence on $\mathcal{O}_K$. Any such norm on $\mathcal{O}_K$ as a vector space over $\mathbf{Q}$ can be used as Size.

1.1 Correlation functions

The first step is the construction of correlation functions for $\mathcal{F}_m$. Choose r integers $a_1, \ldots, a_r \in \mathcal{O}_K$ and consider the following subset of $\mathcal{O}_K$:

$$\mathcal{F}_m^x(a_1, a_2, \ldots, a_r) = \{a \in B_x : \ \mu_m(a) = \mu_m(a + a_1) = \ldots = \mu_m(a + a_r) = 1\}.$$

The ratio

$$\mathbf{E}_m^x(a_1, a_2, \ldots, a_r) := \frac{\#\mathcal{F}_m^x(a_1, a_2, \ldots, a_r)}{\#B_x}$$

is the frequency of square-free integers $a \in \mathcal{O}_K$ for which $a + a_1, a + a_2, \ldots, a + a_r$ are also m-free. It also gives the expectation (hence the notation $\mathbf{E}$) of the product $\mu_m(a)\mu_m(a + a_1) \cdots \mu_m(a + a_r)$ with respect to the uniform measure on B_x. We prove the

Proposition 1.1. *For every $r \geqslant 1$ and every $a_1, a_2, \ldots, a_r \in \mathcal{O}_K$, the limit*

$$c_{r+1}(a_1, \ldots, a_r) = \lim_{x \to \infty} \mathbf{E}_m^x(a_1, \ldots, a_r) \tag{2}$$

exists and

$$c_{r+1}(a_1, \ldots, a_r) = \prod_{\mathfrak{p}} \left(1 - \frac{D(\mathfrak{p}^m \mid 0, a_1, \ldots, a_r)}{N(\mathfrak{p}^m)}\right), \tag{3}$$

where

$$D(\mathfrak{a} \mid a_1, \ldots, a_s) = \#\{b \mod \mathfrak{a} : \ b \equiv a_i \mod \mathfrak{a} \ \textit{for some } i\}.$$

We shall refer to c_{r+1} as the *$(r+1)$-st correlation function for $\mathcal{F}_m$*; we will not indicate the dependence on m explicitly. Notice that, if $\pi_{\mathfrak{a}} : \mathcal{O}_K \to \mathcal{O}_K/\mathfrak{a}$ is the natural projection, then $D(\mathfrak{p}^m \mid 0, a_1, \ldots, a_r) = \#\pi_{\mathfrak{p}^m}(\{0, a_1, a_2, \ldots, a_r\})$ counts the distinct cosets modulo $\mathfrak{p}^m$.

Observe that $\mathcal{F}_m^x(0)$ is simply the set of all m-free integers a such that $\operatorname{Size} a \leqslant x$. By taking $r = 1$ and $a_1 = 0$ in Proposition 1.1, we have the well known

Corollary 1.2. *The asymptotic density of m-free integers in $\mathcal{O}_K$ is*

$$c_2(0) = \lim_{x \to \infty} \mathbf{E}_m^x(0) = \lim_{x \to \infty} \frac{1}{\#B_x} \sum_{a \in B_x} \mu_m(a) = \prod_{\mathfrak{p}} \left(1 - \frac{1}{N(\mathfrak{p}^m)}\right) = \frac{1}{\zeta_K(m)}. \tag{4}$$

Proposition 1.1 is a particular case of Proposition 2.9 given in Section 2. For fixed $a_1, \ldots, a_r \in \mathcal{O}_K$, we have $D(\mathfrak{p}^m \mid 0, a_1, \ldots, a_r) = r + 1$ for all but finitely many prime ideals $\mathfrak{p}$, for which $1 \leqslant D(\mathfrak{p}^m \mid 0, a_1, \ldots, a_r) \leqslant r$. Moreover, let us observe that some correlations are zero (see e.g. Section 6).

For $K = \mathbf{Q}$, the study of $\mathbf{E}_m^x(a_1, \ldots, a_r)$ as $x \to \infty$ is classical, see L. Mirsky [8], R.R. Hall [4], K.M. Tsang [15], D.R. Heath-Brown [6]. There is a geometric notion of m-freeness for points in an arbitrary lattice, studied by M. Baake, R.V. Moody and P.A.B. Pleasants [1] and by Pleasants and C. Huck [11], for which the second correlation function (along with entropies and diffraction spectra) has been computed explicitly. This notion of m-freeness agrees with the one discussed above only when $K = \mathbf{Q}$.

1.1.1 The Spectral Measure and the m-Free Solenoid

For an ideal $\mathfrak{a} \subseteq \mathcal{O}_K$, let us denote by $\Lambda_{\mathfrak{a}}$ the set of unitary characters $\chi : \mathcal{O}_K \to \mathbf{S}^1$ such that $\chi(a) = 1$ for all $a \in \mathfrak{a}$. Notice that $\#\Lambda_{\mathfrak{a}} = \#\mathcal{O}_K/\mathfrak{a} = N(\mathfrak{a})$.

Lemma 1.3. *Let us consider the measure*

$$\nu = \sum_{\mu^2(\mathfrak{d})=1} \sigma_{\mathfrak{d}} \sum_{\chi \in \Lambda_{\mathfrak{d}^m}} \delta_\chi$$

on $\widehat{\mathcal{O}_K}$*, where*

$$\sigma_{\mathfrak{d}} = \sum_{\substack{\mathfrak{b}_0, \mathfrak{b}_1 \subseteq \mathcal{O}_K \\ \mu_m(\mathfrak{b}_0) = \mu_m(\mathfrak{b}_1) = 1 \\ \gcd(\mathfrak{b}_0, \mathfrak{b}_1) = \mathfrak{d}}} \frac{\mu(\mathfrak{b}_0)\mu(\mathfrak{b}_1)}{N(\operatorname{lcm}(\mathfrak{b}_0, \mathfrak{b}_1)^m)}. \tag{5}$$

Then $\widehat{\nu}(a) = c_2(a)$*,* $a \in \mathcal{O}_K$*.*

We shall refer to ν as the *spectral measure.* Let Λ be the support of ν. It is automatically a group and, by the Chinese Remainder Theorem for ideals,

$$\Lambda = \bigcup_{\mu^2(\mathfrak{d})=1} \Lambda_{\mathfrak{d}^m} \cong \bigoplus_{\mathfrak{p}} \mathcal{O}_K/\mathfrak{p}^m.$$

By Pontryagin duality, $\widehat{\Lambda}$ is isomorphic to the compact abelian group

$$\mathbf{G} = \prod_{\mathfrak{p}} \mathcal{O}_K/\mathfrak{p}^m, \tag{6}$$

called the *m-free solenoid.* Elements of $\mathbf{G}$ are coset sequences indexed by the set of prime ideals in $\mathcal{O}_K$, i.e. $\mathbf{g} = (g_{\mathfrak{p}_1^m} + \mathfrak{p}_1^m, g_{\mathfrak{p}_2^m} + \mathfrak{p}_2^m, \ldots) = (g_{\mathfrak{p}^m} + \mathfrak{p}^m)_{\mathfrak{p}}$, where $g_{\mathfrak{p}^m} + \mathfrak{p}^m \in \mathcal{O}_K/\mathfrak{p}^m$. Given $\mathbf{h} \in \mathbf{G}$, we denote by $\mathbf{T_h}$ the translation $\mathbf{T_h}(\mathbf{g}) = \mathbf{g} + \mathbf{h}$. The Haar measure on $\mathbf{G}$ is the product of the counting measures on each factor $\mathcal{O}_K/\mathfrak{p}^m$ and is defined on the natural Borel σ-algebra on $\mathbf{G}$.

We have a $\mathbf{Z}^d$-action on $\mathbf{Z}^d \curvearrowright (\mathbf{G}, \text{Haar})$ as follows: if $\mathbf{Z}^d \ni v$ and $\mathbf{G} \ni \mathbf{g} = (g_{\mathfrak{p}^m} + \mathfrak{p}^m)_{\mathfrak{p}}$, then

$$v \cdot \mathbf{g} = \left([g_{\mathfrak{p}^m} + \iota(v)]\right)_{\mathfrak{p}}. \tag{7}$$

In other words, $\mathbf{Z}^d$ acts by d commuting translations $\mathbf{T}_{\mathbf{u}_1}, \ldots, \mathbf{T}_{\mathbf{u}_d}$ on $\mathbf{G}$, where $\mathbf{u}_i = (e_i + \mathfrak{p}^m)_{\mathfrak{p}} \in \mathbf{G}$.

1.2 A Natural $\mathcal{O}_K$-Action and Main Theorem

Consider the space $X = \{0,1\}^{\mathcal{O}_K}$, whose elements are $\mathcal{O}_K$-indexed sequences $x = (x(a))_{a\in\mathcal{O}_K}$, equipped with the Borel σ-algebra generated by cylinder sets. Introduce on X the probability measure Π defined as follows: for every $r \geqslant 0$ and every $a_0, a_1, \ldots, a_r \in \mathcal{O}_K$

$$\begin{aligned}&\Pi\left\{x \in X:\ x(a_0) = x(a_1) = \ldots = x(a_r) = 1\right\} =\\ &= \zeta_K(m)\cdot c_{r+1}(a_1 - a_0, a_2 - a_0, \ldots, a_r - a_0),\end{aligned} \tag{8}$$

where c_{r+1} is the $(r+1)$-st correlation function (3) associated to $\mathcal{F}_m$. It is clear that (8) determines the measure Π uniquely. We call Π *the natural measure corresponding to the set of m-free integers in $\mathcal{O}_K$.*

If we consider the $\mathcal{O}_K$-action on X defined as $b\cdot x = (x(a+b))_{a\in\mathcal{O}_K}$ for $b \in \mathcal{O}_K$ and $x \in X$, then it follows immediately from (8) that Π is invariant under this action. We can now formulate the main result of this paper.

Main Theorem. *(i) The action $\mathcal{O}_K \curvearrowright (X,\Pi)$ is ergodic and has pure point spectrum given by Λ.*

(ii) The two actions $\mathcal{O}_K \curvearrowright (X,\Pi)$ and $\mathbf{Z}^d \curvearrowright (\mathbf{G}, \mathrm{Haar})$ given in (7) are isomorphic.

Corollary 1.4. *The action $\mathcal{O}_K \curvearrowright (X,\Pi)$ is not weakly mixing and it has entropy zero.*

The above results generalize the work of the fist author and Ya.G. Sinai [3], in which $K = \mathbf{Q}$ and $m = 2$. In this case Corollary 1.4 was also proven by P. Sarnak [14].

The paper is organized as follows. Section 2 contains a generalization of the work by Mirsky [8] to the case of m-free ideals. This implies, in particular, (2), (3) and (4). Section 3 provides other useful formulæ for correlations and two useful lemmata concerning averages and character sums for correlation functions. These results are crucial for the spectral analysis of the action $\mathcal{O}_K \curvearrowright (X,\Pi)$. This analysis is carried out in Section 4 and yields the first part of Main Theorem. In Section 5 we analyze the spectrum of the action of $\mathbf{Z}^d$ on $(\mathbf{G}, \mathrm{Haar})$ and use Mackey's isomorphism theorem for pure point actions [7] to prove the second part of Main Theorem. Section 6 discusses the example of $K = \mathbf{Q}(i)$ where $\mathcal{O}_K = \mathbf{Z}[i]$ is the ring of Gaussian integers.

Acknowledgments

We would like to thank Yakov G. Sinai for his encouragement and advice, Ali Altuğ for helpful suggestions, and M. Baake for bringing the references [1, 11] to our attention. Special appreciation goes to Idris Assani, organizer of the *Ergodic Theory Workshop* in Chapel Hill NC, where part of this work was carried out. The second author acknowledges the support of the ERC Advanced Grant HFAKT.

2 Arithmetical Pattern Problems for m-Free Ideals

Fix an integer $m \geqslant 2$, and recall that μ_m is the indicator of m-free ideals. Define the function D by

$$D(\mathfrak{a} \mid a_1, \ldots, a_s) = \#\{b \bmod \mathfrak{a} : \ b \equiv a_i \bmod \mathfrak{a} \text{ for some } i\}.$$

More generally set

$$D(\mathfrak{a}, \mathfrak{b} \mid a_1, \ldots, a_s) = \#\{b \bmod \mathfrak{a} \mid b \equiv a_i \bmod \mathfrak{a} \text{ for some } i \text{ and } b \in \gcd(\mathfrak{a}, \mathfrak{b})\}.$$

Also set

$$M_m(x; \mathfrak{b}; a_1, \ldots, a_s) = \sum_{\substack{a \in \mathfrak{b} \\ \operatorname{Size} a \leqslant x}} \mu_m(a + a_1) \ldots \mu_m(a + a_s).$$

For ideals $\mathfrak{n}_1, \ldots, \mathfrak{n}_s$ define the E symbol by

$$E\begin{pmatrix} \mathfrak{n}_1 & \ldots & \mathfrak{n}_s \\ a_1 & \ldots & a_s \end{pmatrix} = \begin{cases} 1 & \text{if } \exists b \text{ s.t. } b + a_i \equiv 0 \mod \mathfrak{n}_i \, \forall i \\ 0 & \text{otherwise.} \end{cases} \tag{9}$$

Lemma 2.1. *Equation* (9) *evaluates to 1 iff*

$$a_i - a_j \equiv 0 \mod \gcd(\mathfrak{n}_i, \mathfrak{n}_j) \, \forall i, j.$$

In this case there is exactly one b in each residue class modulo $\operatorname{lcm}(\mathfrak{n}_1, \ldots, \mathfrak{n}_s)$.

Proof. It is enough to observe that

$$\begin{gathered} \mathcal{O}_K / \operatorname{lcm}(\mathfrak{n}_1, \ldots, \mathfrak{n}_s) \to \mathcal{O}_K/\mathfrak{n}_1 \times \cdots \times \mathcal{O}_K/\mathfrak{n}_s \\ a + \operatorname{lcm}(\mathfrak{n}_1, \ldots, \mathfrak{n}_s) \mapsto (a + \mathfrak{n}_1, \ldots, a + \mathfrak{n}_s) \end{gathered}$$

is an isomorphism onto its image. □

Lemma 2.2. *Let*

$$T(x) = \#\{b \in \mathcal{O}_K \mid \operatorname{Size} b \leqslant x \text{ and } b + a_i \equiv 0 \mod \mathfrak{n}_i \, \forall i\}$$

with notation as before. Then

$$\left| T(x) - x^d \frac{E\begin{pmatrix} \mathfrak{n}_1 & \ldots & \mathfrak{n}_s \\ a_1 & \ldots & a_s \end{pmatrix}}{N(\operatorname{lcm}(\mathfrak{n}_1, \ldots, \mathfrak{n}_s))} \right| = O\left(x^{d-1} \operatorname{diam} \operatorname{lcm}(\mathfrak{n}_1, \ldots, \mathfrak{n}_s)\right).$$

Proof. We omit the proof as it is standard.

□

Lemma 2.3. *Let* $T(x)$ *be the number of solutions to the system*

$$\begin{cases} b_i - b = a_i \text{ for all } 1 \leqslant i \leqslant s \\ \quad b_i \in \mathfrak{n}_i \\ \quad b \in \mathfrak{b} \end{cases}$$

such that $\operatorname{Size} b \leqslant x$. *Then we have*

$$\left| T(x) - x^d \frac{E\begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1 & \dots & \mathfrak{n}_s \\ 0 & a_1 & \dots & a_s \end{pmatrix}}{N(\operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1, \dots, \mathfrak{n}_s))} \right| = O\left(x^{d-1} \operatorname{diam} \operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1, \dots, \mathfrak{n}_s)\right) \tag{10}$$

Proof. Apply Lemmata 2.1 and 2.2 with s replaced by $s+1$. $\square$

Lemma 2.4. *If a function is multiplicative on ideals, then it is determined by its values at prime powers. That is, if*

$$\sum_{\mathfrak{n}_1, \dots, \mathfrak{n}_s} |f(\mathfrak{n}_1, \dots, \mathfrak{n}_s)| < \infty$$

and

$$f(\mathfrak{n}_1, \dots, \mathfrak{n}_s) f(\mathfrak{n}'_1, \dots, \mathfrak{n}'_s) = f(\mathfrak{n}_1\mathfrak{n}'_1, \dots, \mathfrak{n}_s\mathfrak{n}'_s) \text{ whenever } \gcd(\mathfrak{n}_i, \mathfrak{n}'_j) = \mathcal{O}_K \text{ for all } i, j,$$

then

$$\sum_{\mathfrak{n}_1, \dots, \mathfrak{n}_s} f(\mathfrak{n}_1, \dots, \mathfrak{n}_s) = \prod_{\mathfrak{p}} \chi_{\mathfrak{p}}$$

where

$$\chi_{\mathfrak{p}} = \sum_{\delta_1, \dots, \delta_s \geqslant 0} f(\mathfrak{p}^{\delta_1}, \dots, \mathfrak{p}^{\delta_s}).$$

Proof. Follows by a simple induction. $\square$

Lemma 2.5. *Take* $t \geqslant 1$ *and* $m \geqslant 2$. *Also fix a prime ideal* $\mathfrak{p}$ *and* $k_1, \dots, k_t \in \mathcal{O}_K$ *such that*

$$k_1 \equiv \dots \equiv k_t \equiv \rho \mod \mathfrak{p}^m.$$

Then

$$\sum_{\substack{\eta_1, \dots, \eta_t \in \{0,1\} \\ \text{not all zero}}} (-1)^{\eta_1 + \dots + \eta_t} E\begin{pmatrix} \mathfrak{b} & \mathfrak{p}^{m\eta_1} & \dots & \mathfrak{p}^{m\eta_t} \\ 0 & k_1 & \dots & k_t \end{pmatrix} = \begin{cases} -1 & \text{if } \rho \in \gcd(\mathfrak{p}^m, \mathfrak{b}) \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Note that η_i is non-zero for at least one i, so that

$$E\begin{pmatrix} \mathfrak{b} & \mathfrak{p}^{m\eta_1} & \dots & \mathfrak{p}^{m\eta_t} \\ 0 & k_1 & \dots & k_t \end{pmatrix} = E\begin{pmatrix} \mathfrak{b} & \mathfrak{p}^m \\ 0 & \rho \end{pmatrix} = \begin{cases} 1 & \rho \in \gcd(\mathfrak{p}^m, \mathfrak{b}) \\ 0 & \text{otherwise} \end{cases}$$

by Lemma 2.1. Furthermore

$$\sum_{\substack{\eta_1,\dots,\eta_t \in \{0,1\} \\ \text{not all zero}}} (-1)^{\eta_1+\dots+\eta_t} = -1,$$

whence the result. ☐

Lemma 2.6. *With notation as before set*

$$S = S_{m,\mathfrak{b}}(a_1,\dots,a_s) = \sum_{\mathfrak{n}_1,\dots,\mathfrak{n}_s} \frac{\mu(\mathfrak{n}_1)\dots\mu(\mathfrak{n}_s)}{N(\operatorname{lcm}(\mathfrak{b},\mathfrak{n}_1^m,\dots,\mathfrak{n}_s^m))} E\begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1^m & \dots & \mathfrak{n}_s^m \\ 0 & a_1 & \dots & a_s \end{pmatrix}.$$

Then we have

$$S = \frac{1}{N(\mathfrak{b})} \prod_{\mathfrak{p}^m \not\supseteq \mathfrak{b}} \left(1 - \frac{N(\gcd(\mathfrak{p}^m,\mathfrak{b}))D(\mathfrak{p}^m,\mathfrak{b} \mid a_1,\dots,a_s)}{N(\mathfrak{p}^m)}\right),$$

and it vanishes precisely when

$$D(\mathfrak{p}^m,\mathfrak{b} \mid a_1,\dots,a_s) = \frac{N(\mathfrak{p}^m)}{N(\gcd(\mathfrak{p}^m,\mathfrak{b}))} \text{ for some } \mathfrak{p}.$$

Proof. We claim that if

$$\gcd(\mathfrak{n}_i,\mathfrak{n}_j') = \mathcal{O}_K \tag{11}$$

for all i and j, then

$$\frac{1}{N(\mathfrak{b})}N(\operatorname{lcm}(\mathfrak{b},\mathfrak{n}_1,\dots,\mathfrak{n}_s)) \cdot \frac{1}{N(\mathfrak{b})}N(\operatorname{lcm}(\mathfrak{b},\mathfrak{n}_1',\dots,\mathfrak{n}_s')) = \frac{1}{N(\mathfrak{b})}N(\operatorname{lcm}(\mathfrak{b},\mathfrak{n}_1\mathfrak{n}_1',\dots,\mathfrak{n}_s\mathfrak{n}_s'))$$

and

$$E\begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1 & \dots & \mathfrak{n}_s \\ 0 & a_1 & \dots & a_s \end{pmatrix} \cdot E\begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1' & \dots & \mathfrak{n}_s' \\ 0 & a_1 & \dots & a_s \end{pmatrix} = E\begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1\mathfrak{n}_1' & \dots & \mathfrak{n}_s\mathfrak{n}_s' \\ 0 & a_1 & \dots & a_s \end{pmatrix}. \tag{12}$$

For each prime ideal $\mathfrak{p}$ let $\mathfrak{p}^\lambda$, $\mathfrak{p}^{\nu_i}$, $\mathfrak{p}^{\nu_j'}$ be its largest powers that divide $\mathfrak{b}$, $\mathfrak{n}_i$, and $\mathfrak{n}_j'$, respectively. Then it is enough to confirm that

$$\max(\lambda,\nu_1,\dots,\nu_s) + \max(\lambda,\nu_1',\dots,\nu_s') - 2\lambda = \max(\lambda,\nu_1+\nu_1',\dots,\nu_s+\nu_s') - \lambda.$$

From (11) it follows that $\nu_i = 0$ for all i or $\nu_j' = 0$ for all j, so the preceding equation is verified.

For the second claim (12) note that if the left-hand side vanishes, then so does the right-hand side. So suppose the left-hand side doesn't vanish, that is,

$$\begin{aligned} a_i - a_j &\in \gcd(\mathfrak{n}_i, \mathfrak{n}_j), \quad a_i \in \gcd(\mathfrak{b}, \mathfrak{n}_i) \\ a_i - a_j &\in \gcd(\mathfrak{n}_i', \mathfrak{n}_j'), \quad a_i \in \gcd(\mathfrak{b}, \mathfrak{n}_i') \end{aligned}$$

by Lemma 2.1. From (11) these conditions are equivalent to

$$a_i - a_j \in \gcd(\mathfrak{n}_i\mathfrak{n}_i', \mathfrak{n}_j\mathfrak{n}_j'), \quad a_i \in \gcd(\mathfrak{b}, \mathfrak{n}_i\mathfrak{n}_i'),$$

and the claim follows by another application of Lemma 2.1.

Now then we write

$$N(\mathfrak{b})S = \sum_{\mathfrak{n}_1,\dots,\mathfrak{n}_s} \frac{\mu(\mathfrak{n}_1)\dots\mu(\mathfrak{n}_s)N(\mathfrak{b})}{N(\operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1^m, \dots, \mathfrak{n}_s^m))} E\begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1^m & \dots & \mathfrak{n}_s^m \\ 0 & c_1 & \dots & c_s \end{pmatrix} = \sum_{\mathfrak{n}_1,\dots,\mathfrak{n}_s} f(\mathfrak{n}_1, \dots, \mathfrak{n}_s).$$

This sum converges absolutely

$$\begin{aligned} \sum_{\mathfrak{n}_1,\dots,\mathfrak{n}_s} |f(\mathfrak{n}_1, \dots, \mathfrak{n}_s)| &\leqslant N(\mathfrak{b}) \sum \frac{1}{N(\operatorname{lcm}(\mathfrak{n}_1^m, \dots, \mathfrak{n}_s^m))} \\ &\ll \sum_{\mathfrak{n}} \frac{1}{N^m(\mathfrak{n})} \sum_{\operatorname{lcm}(\mathfrak{n}_i)=\mathfrak{n}} 1 \ll \sum_{\mathfrak{n}} \frac{d^s(\mathfrak{n})}{N^m(\mathfrak{n})} \ll \zeta_K(m-\varepsilon) < \infty \end{aligned}$$

since $m \geqslant 2$ and the divisor function satisfies $d(\mathfrak{n}) \ll_\varepsilon N^\varepsilon(\mathfrak{n})$. By multiplicativity and Lemma 2.4 we have

$$N(\mathfrak{b})S = \prod_{\mathfrak{p}} \chi_{\mathfrak{p}},$$

with

$$\begin{aligned} \chi_{\mathfrak{p}} &= \sum_{\delta_1,\dots,\delta_s \geqslant 0} \frac{(-1)^{\delta_1+\dots+\delta_s} N(\mathfrak{b})}{N(\operatorname{lcm}(\mathfrak{b}, \mathfrak{p}^{m\delta_1}, \dots, \mathfrak{p}^{m\delta_s}))} E\begin{pmatrix} \mathfrak{b} & \mathfrak{p}^{m\delta_1} & \dots & \mathfrak{p}^{m\delta_s} \\ 0 & a_1 & \dots & a_s \end{pmatrix} = \\ &= 1 + \frac{N(\gcd(\mathfrak{p}^m, \mathfrak{b}))}{N(\mathfrak{p}^m)} \psi_{\mathfrak{p}} \end{aligned}$$

and

$$\psi_{\mathfrak{p}} = \sum_{\substack{\delta_1,\dots,\delta_s \in \{0,1\} \\ \text{not all zero}}} (-1)^{\delta_1+\dots+\delta_s} E\begin{pmatrix} \mathfrak{b} & \mathfrak{p}^{m\delta_1} & \dots & \mathfrak{p}^{m\delta_s} \\ 0 & a_1 & \dots & a_s \end{pmatrix}. \tag{13}$$

It remains to evaluate $\psi_{\mathfrak{p}}$. Observe that the terms of (13) are zero if a_j are not congruent modulo $\mathfrak{p}^m$ for all $j \in \{i : \delta_i = 1\}$. For ρ modulo $\mathfrak{p}^m$ let t_ρ denote the number of integers

$a_1, \dots, a_s$ that are congruent to ρ. If $t_\rho > 0$, denote them by $k_1^{(\rho)}, \dots, k_{t_\rho}^{(\rho)}$. Then using Lemma 2.5 we have

$$\begin{aligned}\psi_{\mathfrak{p}} &= \sum_{\substack{\rho \mod \mathfrak{p}^m \\ t_\rho > 0}} \sum_{\substack{\eta_1, \dots, \eta_{t_\rho} \in \{0,1\} \\ \text{not all zero}}} (-1)^{\eta_1 + \dots + \eta_{t_\rho}} E \begin{pmatrix} \mathfrak{b} & \mathfrak{p}^{m\eta_1} & \dots & \mathfrak{p}^{m\eta_{t_\rho}} \\ 0 & k_1^{(\rho)} & \dots & k_{t_\rho}^{(\rho)} \end{pmatrix} = \\ &= - \sum_{\substack{\rho \mod \mathfrak{p}^m \\ t_\rho > 0 \\ \rho \in \gcd(\mathfrak{p}^m, \mathfrak{b})}} 1 = -D(\mathfrak{p}^m, \mathfrak{b} \mid a_1, \dots, a_s).\end{aligned}$$

Thus we have

$$N(\mathfrak{b}) S = \prod_{\mathfrak{p}} \left(1 - \frac{N(\gcd(\mathfrak{p}^m, \mathfrak{b}))}{N(\mathfrak{p}^m)} D(\mathfrak{p}^m, \mathfrak{b} \mid a_1, \dots, a_s) \right).$$

It remains to verify the positivity part. Clearly the product vanishes if one of the factors does. Otherwise the factors cannot be less than $1 - \frac{s}{N^m(\mathfrak{p})}$ for $N(\mathfrak{p})$ large enough, and these give a non-zero product. $\square$

Lemma 2.7. *For all x, y in $\mathcal{O}_K$ we have*

$$\operatorname{Size}(xy) \ll \operatorname{Size}(x) \operatorname{Size}(y)$$

and the implied constant depends on ι.

Proof. Let $\{e_i\}_{i=1}^d$ be the basis used to define ι. Then $e_i e_j = \sum_k c_k^{ij} e_k$ for certain $c_k^{ij} \in \mathcal{O}_K$. We have

$$\begin{aligned}\operatorname{Size}(xy) &= \operatorname{Size}\left(\sum_i x_i e_i \sum_j y_j e_j \right) = \operatorname{Size}\left(\sum_{i,j,k} x_i y_j c_k^{ij} e_k \right) \leqslant \\ &\leqslant \max |c_k^{ij}| d \sum_{i,j} |x_i y_j| \ll \left(\sum_i |x_i| \right) \left(\sum_j |y_j| \right) \ll \operatorname{Size}(x) \operatorname{Size}(y).\end{aligned}$$

$\square$

Lemma 2.8. *For every $a \in \mathcal{O}_K \smallsetminus \{0\}$ we have $N(a) \ll \operatorname{Size}^d(a)$, and the implied constant depends on the choice of the isomorphism ι.*

Proof. Fix $b \in \mathcal{O}_K$ whose irreducible polynomial has degree d (it exists by the Primitive Element Theorem). Let $\Xi = \mathbf{Z}[a, ab, ab^2, \dots, ab^{d-1}]$. Observe that $\Xi < \mathcal{O}_K$ is a finite index additive subgroup and $\Xi \subset (a)$, implying that $\# \mathcal{O}_K / \Xi \geqslant N((a))$. Using Lemma 2.7 we have

$$N(a) \leqslant \# \mathcal{O}_K / \Xi \leqslant \prod_{i=0}^{d-1} \operatorname{Size}(ab^i) \ll \operatorname{Size}^d(a) \prod_{i=0}^{d-1} \operatorname{Size}(b^i) \ll \operatorname{Size}^d(a),$$

as needed. Note that minimizing $\prod_{i=0}^{d-1} \operatorname{Size}(b^i)$ over b with full degree irreducible polynomial will remove the dependence on the choice of b. $\square$

Proposition 2.9. *If* $D(\mathfrak{p}^m, \mathfrak{b} \mid a_1, \dots, a_s) = \frac{N(\mathfrak{p}^m)}{N(\gcd(\mathfrak{p}^m, \mathfrak{b}))}$ *for some* $\mathfrak{p}$*, then*

$$M_m(x; \mathfrak{b}; a_1, \dots, a_s) = 0.$$

Otherwise

$$M_m(x; \mathfrak{b}; a_1, \dots, a_s) = S_{m,\mathfrak{b}}(a_1, \dots, a_s) x^d + O(x^{d - \frac{m-1}{m+2sm-1}})$$

and the implied constant depends on ι*.*

Proof. Say $D(\mathfrak{p}^m, \mathfrak{b} \mid a_1, \dots, a_s) = \frac{N(\mathfrak{p}^m)}{N(\gcd(\mathfrak{p}^m, \mathfrak{b}))}$ for some $\mathfrak{p}$. Then it is easy to see that $a + a_i \in \mathfrak{p}^m$ for some i, proving the first case.

Suppose then that $D(\mathfrak{p}^m, \mathfrak{b} \mid a_1, \dots, a_s) < \frac{N(\mathfrak{p}^m)}{N(\gcd(\mathfrak{p}^m, \mathfrak{b}))}$ for all $\mathfrak{p}$. Let $\alpha \in (0, 1/m)$ to be chosen later. From the relation (1) we get that

$$\begin{aligned} M_m(x; \mathfrak{b}; a_1, \dots, a_s) &= \sum_{\substack{\text{Size}\, a \leqslant x \\ a \in \mathfrak{b}}} \sum_{\substack{a + c_i \in \mathfrak{n}_i^m \\ 1 \leqslant i \leqslant s}} \mu(\mathfrak{n}_1) \dots \mu(\mathfrak{n}_s) = \sum_{\mathfrak{n}_i} \mu(\mathfrak{n}_1) \dots \mu(\mathfrak{n}_s) \sum_{\substack{\text{Size}\, a \leqslant x \\ a \in \mathfrak{b} \\ a + c_i \in \mathfrak{n}_i^m}} 1 = \\ &= \Sigma_1 + \Sigma_2. \end{aligned}$$

The first sum is over s-tuples of ideals $\mathfrak{n}_i$ of norm at most x^α, while Σ_2 includes s-tuples where at least one ideal has norm greater than x^α. By Lemma 2.3,

$$\begin{aligned} &\Sigma_1 = \\ &= \sum_{N(\mathfrak{n}_i) \leqslant x^\alpha} \mu(\mathfrak{n}_1) \dots \mu(\mathfrak{n}_s) \left(x^d \frac{E \begin{pmatrix} \mathfrak{b} & \mathfrak{n}_1^m & \dots & \mathfrak{n}_s^m \\ 0 & a_1 & \dots & a_s \end{pmatrix}}{N(\operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1^m, \dots, \mathfrak{n}_s^m))} + O(x^{d-1} \operatorname{diam} \operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1^m, \dots, \mathfrak{n}_s^m)) \right) = \\ &= \frac{x^d}{N(\mathfrak{b})} \prod_{\mathfrak{p}^m \not\supseteq \mathfrak{b}} \left(1 - \frac{N(\gcd(\mathfrak{p}^m, \mathfrak{b})) D(\mathfrak{p}^m, \mathfrak{b} \mid a_1, \dots, a_s)}{N(\mathfrak{p}^m)} \right) + \\ &+ O\left(x^d \sum_{k=1}^{s} \sum_{\substack{N(\mathfrak{n}_k) > x^\alpha \\ \mathfrak{n}_1, \dots, \mathfrak{n}_s}} \frac{1}{N(\operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1^m, \dots, \mathfrak{n}_s^m))} \right) + O\left(x^{d-1} \sum_{N(\mathfrak{n}_i) \leqslant x^\alpha} \operatorname{diam} \operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1^m, \dots, \mathfrak{n}_s^m) \right). \end{aligned} \tag{14}$$

The first error term from (14) is

$$\begin{aligned} &\ll x^d \sum_{N(\mathfrak{n}) > x^\alpha} \frac{d^s(\mathfrak{n})}{N^m(\mathfrak{n})} \ll_\varepsilon \sum N^{-m+\varepsilon}(\mathfrak{n}) \left(\frac{N(\mathfrak{n})}{x^\alpha} \right)^{m-1-2\varepsilon} \ll \\ &\ll x^{d + \alpha(1 - m + 2\varepsilon)} \zeta_K(1 + \varepsilon) \ll x^{d - \alpha(m-1) + \varepsilon}. \end{aligned} \tag{15}$$

The second error term is bounded by

$$\begin{aligned} &x^{d-1} \sum_{N(\mathfrak{n}_i) \leqslant x^\alpha} N(\operatorname{lcm}(\mathfrak{b}, \mathfrak{n}_1^m, \ldots, \mathfrak{n}_s^m)) \ll x^{d-1} \sum_{N(\mathfrak{n}) < x^{\alpha s m}} N(\mathfrak{n}) d^s(\mathfrak{n}) \ll \\ &\ll x^{d-1} \sum N^{1+\varepsilon}(\mathfrak{n}) \left(\frac{x^{\alpha s m}}{N(\mathfrak{n})} \right)^{2+2\varepsilon} \ll x^{d-1+\alpha s m(2+2\varepsilon)} \zeta_K(1+\varepsilon) \ll x^{d-1+2\alpha s m+\varepsilon}. \quad (16) \end{aligned}$$

For Σ_2 we have

$$|\Sigma_2| \leqslant \sum_{k=1}^{s} f_k$$

and

$$\begin{aligned} |f_k| \leqslant &\sum_{\substack{\text{Size}\, a \leqslant x \\ a+a_i \in \mathfrak{n}_i^m \\ a \in \mathfrak{b} \\ N(\mathfrak{n}_k) > x^\alpha}} 1 = \sum_{\substack{a+a_k \in \mathfrak{n}_k^m \\ \text{Size}\, a \leqslant x \\ N(\mathfrak{n}_k) > x^\alpha}} \prod_{i \neq k} \sum_{a+a_i \in \mathfrak{n}_i^m} 1 \leqslant \sum \prod d((a+a_i)) \ll_\varepsilon \sum \prod N^\varepsilon((a+a_i)) \ll \\ &\ll x^\varepsilon \sum_{\substack{a+a_k \in \mathfrak{n}_k^m \\ \text{Size}\, a \leqslant x \\ N(\mathfrak{n}_k) > x^\alpha}} 1 \ll x^\varepsilon \sum_{\substack{\mathfrak{m} \subset \mathfrak{n}^k \\ N(\mathfrak{m}) \ll x^d \\ N(\mathfrak{n}) > x^\alpha}} 1 \ll \quad (17) \\ &\ll \sum_{N(\mathfrak{n}) > x^\alpha} \left(\frac{x^d}{N(\mathfrak{n}^m)} \right)^{1+\varepsilon} \ll x^{d+d\varepsilon} \sum_{N(\mathfrak{n}) > x^\alpha} \frac{1}{N^{m+m\varepsilon}(\mathfrak{n})} \left(\frac{N(\mathfrak{n})}{x^\alpha} \right)^{m+m\varepsilon-1-\varepsilon} \ll \\ &\ll x^{d-\alpha(m-1)+\varepsilon}. \quad (18) \end{aligned}$$

where the inequalities on line (17) are due to Lemma 2.8. The three error terms come from equations (15), (16), and (18); setting them equal gives $\alpha = \frac{1}{m+2sm-1}$ and proves the Proposition. $\square$

Propositon 1.1 in Section 1 is a particular case of Proposition 2.9.

3 More Formulæ for the correlation functions

Recall the definition (5) of $\sigma_{\mathfrak{d}}$ from Section 1.1.1. Throughout the paper, $\mathfrak{d}$ indicates a square-free ideal; equivalently, $\mathfrak{d}$ can be thought as a finite collection of prime ideals (or *places*). We have the

Lemma 3.1.

$$\sigma_{\mathfrak{d}} = \frac{1}{N(\mathfrak{d}^m)} \prod_{\mathfrak{p} \not\supseteq \mathfrak{d}} \left(1 - \frac{2}{N(\mathfrak{p}^m)} \right).$$

Proof. Multiply out the product in the RHS to get the sum 5) defining $\sigma_{\mathfrak{d}}$. $\square$

In particular, Lemma 3.1 shows that $\sigma_{\mathfrak{d}}$ is positive and bounded away from zero and infinity. More precisely

$$0 < \prod_{\mathfrak{p}} \left(1 - \frac{2}{N(\mathfrak{p}^2)}\right) = \sigma_{\mathcal{O}_K} \leqslant \sigma_{\mathfrak{d}} < \frac{1}{\zeta_K(m)}$$

and we can also write

$$\sigma_{\mathfrak{d}} = \sigma_{\mathcal{O}_K} \prod_{\mathfrak{p} \supseteq \mathfrak{d}} \frac{1}{N(\mathfrak{p}^2) - 2}. \tag{19}$$

Lemma 3.2. *Let $a \in \mathcal{O}_K$. Then*

$$c_2(a) = \sum_{\substack{\mathfrak{d}^m \supseteq (a) \\ \mu^2(\mathfrak{d}) = 1}} \sigma_{\mathfrak{d}}, \tag{20}$$

and sum converges absolutely.

Proof. From Proposition 1.1 we get

$$D(\mathfrak{p}^m \mid 0, a) = \begin{cases} 1 & \text{if } \mathfrak{p}^m \supseteq (a); \\ 2 & \text{otherwise.} \end{cases}$$

This gives

$$c_2(a) = \prod_{\mathfrak{p}^m \supseteq (a)} \left(1 - \frac{1}{N(\mathfrak{p}^m)}\right) \prod_{\mathfrak{p}^m \not\supseteq (a)} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right). \tag{21}$$

The sum in the RHS of (20) converges absolutely by (19). By Lemma 3.1 we have

$$\sum_{\substack{\mathfrak{d}^m \supseteq (a) \\ \mu^2(\mathfrak{d}) = 1}} \sigma_{\mathfrak{d}} = \sum_{\substack{\mathfrak{d}^m \supseteq (a) \\ \mu^2(\mathfrak{d}) = 1}} \frac{1}{N(\mathfrak{d}^m)} \prod_{\mathfrak{p} \not\supseteq \mathfrak{d}} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right) =$$

$$= \prod_{\mathfrak{p}} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right) \sum_{\substack{\mathfrak{d}^m \supseteq (a) \\ \mu^2(\mathfrak{d}) = 1}} \frac{1}{N(\mathfrak{d}^m)} \prod_{\mathfrak{p} \supseteq \mathfrak{d}} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right)^{-1} =$$

$$= \prod_{\mathfrak{p}} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right) \sum_{\substack{\mathfrak{d}^m \supseteq (a) \\ \mu^2(\mathfrak{d}) = 1}} \prod_{\mathfrak{p} \supseteq \mathfrak{d}} \frac{1}{N(\mathfrak{p}^m) - 2} =$$

$$= \prod_{\mathfrak{p}} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right) \prod_{\mathfrak{p}^m \supseteq (a)} \left(1 + \frac{1}{N(\mathfrak{p}^m) - 2}\right) =$$

$$= \prod_{\mathfrak{p}^m \supseteq (a)} \left(1 - \frac{1}{N(\mathfrak{p}^m)}\right) \prod_{\mathfrak{p}^m \not\supseteq (a)} \left(1 - \frac{2}{N(\mathfrak{p}^m)}\right) = c_2(a),$$

where the last equality comes from (21). □

Recall that $\Lambda = \bigcup_{\mu^2(\mathfrak{d})=1} \Lambda_{\mathfrak{d}^m}$. Notice that if $\mathfrak{d}_1 \supseteq \mathfrak{d}_2$ then $\Lambda_{\mathfrak{d}_1^m} \subseteq \Lambda_{\mathfrak{d}_2^m}$. Let us define

$$\Lambda'_{\mathfrak{d}^m} = \{\chi \in \Lambda_{\mathfrak{d}^m} :\ \chi \in \Lambda_{\mathfrak{d}'^m} \Rightarrow \mathfrak{d} \supseteq \mathfrak{d}'\}\,. \tag{22}$$

In other words

$$\chi \in \Lambda'_{\mathfrak{d}^m} \iff \mathfrak{d} = \gcd\{\mathfrak{d}' :\ \mu^2(\mathfrak{d}') = 1 \text{ and } \chi \in \Lambda_{\mathfrak{d}'^m}\}. \tag{23}$$

Let us give the proof of Lemma 1.3 from Section 1.1.1.

Proof of Lemma 1.3. Using Lemma 3.2 we can write

$$c_2(a) = \sum_{\mu^2(\mathfrak{d})=1} C_{\mathfrak{d}}(a), \quad \text{where} \quad C_{\mathfrak{d}}(a) = \begin{cases} \sigma_{\mathfrak{d}}, & \text{if } \mathfrak{d}^m \supseteq (a); \\ 0, & \text{otherwise.} \end{cases}$$

The function $C_{\mathfrak{d}}$ is constant (equal to $\sigma_{\mathfrak{d}}$) on the lattice $\mathfrak{d}^m$ and zero elsewhere. This function on $\mathcal{O}_K$ is the Fourier transform of a measure on $\widehat{\mathcal{O}_K}$, given by a sum of Dirac δ-measures at the points in the set $\Lambda_{\mathfrak{d}^m}$, with equal weights equal to $\sigma_{\mathfrak{d}}/N(\mathfrak{d}^m)$. The formula for the spectral measure ν and the lemma is proved. ☐

Let $\eta \in \widehat{\mathcal{O}_K}$ be the trivial character, $\eta(a) = 1$ for every $a \in \mathcal{O}_K$. The following Lemma generalizes a formula by R.R. Hall [4].

Lemma 3.3. *For every $r \geqslant 1$ and every $a_1, \ldots, a_r \in \mathcal{O}_K$ we have*[3]

$$c_{r+1}(a_1 \ldots, a_r) = \sum_{\mathfrak{a}_0} \sum_{\mathfrak{a}_1} \cdots \sum_{\mathfrak{a}_r} g(\mathfrak{a}_0) g(\mathfrak{a}_1) \cdots g(\mathfrak{a}_r) \sum_{\substack{\chi_i \in \Lambda'_{\mathfrak{a}_i^m} \\ 0 \leqslant i \leqslant r \\ \chi_0\chi_1\cdots\chi_r = \eta}} \chi_0(0)\chi_1(a_1)\cdots\chi_r(a_r), \tag{24}$$

where

$$g(\mathfrak{a}) = \frac{\mu(\mathfrak{a})}{\zeta_K(m)} \prod_{\mathfrak{p} \supseteq \mathfrak{a}} \frac{1}{N(\mathfrak{p}^m) - 1}. \tag{25}$$

Proof. Notice that the inner sum in (24) does not exceed

$$\sum_{\substack{\chi_i \in \Lambda_{\mathfrak{a}_i^m} \\ 0 \leqslant i \leqslant r \\ \chi_0\chi_1\cdots\chi_r = \eta}} 1 = \frac{N(\mathfrak{a}_0^m \mathfrak{a}_1^m \cdots \mathfrak{a}_r^m)}{N(\operatorname{lcm}(\mathfrak{a}_0, \mathfrak{a}_1, \ldots, \mathfrak{a}_r)^m)}$$

in absolute value. Moreover, for every ideal $\mathfrak{a}$, $|g(\mathfrak{a})| \leqslant \frac{1}{N(\mathfrak{a}^m)}$ and the series

$$\sum_{\mathfrak{a}_0} \sum_{\mathfrak{a}_1} \cdots \sum_{\mathfrak{a}_r} \frac{1}{N(\operatorname{lcm}(\mathfrak{a}_0, \mathfrak{a}_1, \ldots, \mathfrak{a}_r)^m)}$$

---

[3] Notice that $\chi_0(0) = 1$ in (24) and will it be omitted in proof of Lemma 3.3.

converges absolutely. Let us evaluate the inner sum in (24). Let $\mathfrak{a} = \operatorname{lcm}(\mathfrak{a}_0, \mathfrak{a}_1, \ldots, \mathfrak{a}_r)$. Notice that

$$\frac{1}{N(\mathfrak{a}^m)} \sum_{a \in \mathcal{O}_K/\mathfrak{a}^m} \chi_0(a)\chi_1(a)\cdots\chi_r(a) = \begin{cases} 1 & \text{if } \chi_0\chi_1\cdots\chi_r = \eta; \\ 0 & \text{otherwise.} \end{cases}$$

This allows us to rewrite the inner sum in (24) as

$$\frac{1}{N(\mathfrak{a}^m)} \sum_{a \in \mathcal{O}_K/\mathfrak{a}^m} \prod_{i=0}^{r} \sum_{\chi_i \in \Lambda'_{\mathfrak{a}_i^m}} \chi_i(a_i + a) =$$

$$= \frac{1}{N(\mathfrak{a}^m)} \sum_{a \in \mathcal{O}_K/\mathfrak{a}^m} \prod_{i=0}^{r} \sum_{\substack{\mathfrak{b}_i \supseteq \mathfrak{a}_i \\ \mathfrak{b}_i^m \supseteq (a_i + a)}} \mu\left(\frac{\mathfrak{a}_i}{\mathfrak{b}_i}\right) N(\mathfrak{b}_i^m) =$$

$$= \frac{1}{N(\mathfrak{a}^m)} \sum_{\mathfrak{b}_0 \supseteq \mathfrak{a}_0} \sum_{\mathfrak{b}_1 \supseteq \mathfrak{a}_1} \cdots \sum_{\mathfrak{b}_r \supseteq \mathfrak{a}_r} \mu\left(\frac{\mathfrak{a}_0}{\mathfrak{b}_0}\right) \mu\left(\frac{\mathfrak{a}_1}{\mathfrak{b}_1}\right) \cdots \mu\left(\frac{\mathfrak{a}_r}{\mathfrak{b}_r}\right) N(\mathfrak{b}_0^m \mathfrak{b}_1^m \cdots \mathfrak{b}_r^m) \sum_{\substack{a \in \mathcal{O}_K/\mathfrak{a}^m \\ a \equiv -a_i \bmod \mathfrak{b}_i^m \\ 0 \leqslant i \leqslant r}} 1,$$

where $a_0 = 0$. Observe that

$$\sum_{\substack{a \in \mathcal{O}_K/\mathfrak{a}^m \\ a \equiv -a_i \bmod \mathfrak{b}_i^m \\ 0 \leqslant i \leqslant r}} 1 = E\begin{pmatrix} \mathfrak{b}_0^m & \mathfrak{b}_1^m & \ldots & \mathfrak{b}_r^m \\ 0 & a_1 & \ldots & a_r \end{pmatrix} \frac{N(\mathfrak{a}^m)}{N(\operatorname{lcm}(\mathfrak{b}_0, \mathfrak{b}_1, \ldots, \mathfrak{b}_r)^m)}$$

and thus the inner sum in (24) equals

$$\sum_{\mathfrak{b}_0 \supseteq \mathfrak{a}_0} \sum_{\mathfrak{b}_1 \supseteq \mathfrak{a}_1} \cdots \sum_{\mathfrak{b}_r \supseteq \mathfrak{a}_r} \mu\left(\frac{\mathfrak{a}_0}{\mathfrak{b}_0}\right) \mu\left(\frac{\mathfrak{a}_1}{\mathfrak{b}_1}\right) \cdots \mu\left(\frac{\mathfrak{a}_r}{\mathfrak{b}_r}\right) \frac{N(\mathfrak{b}_0^m \mathfrak{b}_1^m \cdots \mathfrak{b}_r^m)}{N(\operatorname{lcm}(\mathfrak{b}_0, \mathfrak{b}_1, \ldots, \mathfrak{b}_r)^m)} E\begin{pmatrix} \mathfrak{b}_0^m & \mathfrak{b}_1^m & \ldots & \mathfrak{b}_r^m \\ 0 & a_1 & \ldots & a_r \end{pmatrix}. \tag{26}$$

Notice that the $\mathfrak{b}_i$'s are necessarily square-free and thus $\mu(\mathfrak{a}_i/\mathfrak{b}_i) = \mu(\mathfrak{a}_i)\mu(\mathfrak{b}_i)$. Let us also observe that, for $i = 0, \ldots, r$,

$$\sum_{\mathfrak{a}_i \subseteq \mathfrak{b}_i} \mu(\mathfrak{a}_i) g(\mathfrak{a}_i) = \frac{\mu^2(\mathfrak{b}_i)}{N(\mathfrak{b}_i^m)}. \tag{27}$$

To see this, for $\mu^2(\mathfrak{b}_i) = 1$, we can write the LHS of (27) as

$$\frac{1}{\zeta_K(m)} \sum_{\mathfrak{a}_i \subseteq \mathfrak{b}_i} \prod_{\mathfrak{p} \supseteq \mathfrak{a}_i} \left(\frac{1}{N(\mathfrak{p}^m)} + \frac{1}{N(\mathfrak{p}^{2m})} + \frac{1}{N(\mathfrak{p}^{3m})} + \ldots \right) =$$

$$= \frac{1}{\zeta_K(m)} \sum_{\mathfrak{a}_i \subseteq \mathfrak{b}_i} \prod_{\mathfrak{p} \supseteq \mathfrak{b}_i} \left(\frac{1}{N(\mathfrak{p}^m)} + \frac{1}{N(\mathfrak{p}^{2m})} + \ldots \right) \prod_{\substack{\mathfrak{p} \supseteq \mathfrak{a}_i \\ \mathfrak{p} \not\supseteq \mathfrak{b}_i}} \left(\frac{1}{N(\mathfrak{p}^m)} + \frac{1}{N(\mathfrak{p}^{2m})} + \ldots \right) =$$

$$= \frac{1}{\zeta_K(m)} \frac{1}{N(\mathfrak{b}_i^m)} \prod_{\mathfrak{p}\supseteq\mathfrak{b}_i} \left(1+\frac{1}{N(\mathfrak{p}^m)}+\frac{1}{N(\mathfrak{p}^{2m})}+\dots\right) \sum_{\mathfrak{a}_i\subseteq\mathfrak{b}_i} \prod_{\substack{\mathfrak{p}\supseteq\mathfrak{a}_i\\ \mathfrak{p}\not\supseteq\mathfrak{b}_i}} \frac{1}{N(\mathfrak{p}^m)-1} =$$

$$= \frac{1}{\zeta_K(m)} \frac{1}{N(\mathfrak{b}_i^m)} \prod_{\mathfrak{p}\supseteq\mathfrak{b}_i} \left(1-\frac{1}{N(\mathfrak{p}^m)}\right)^{-1} \prod_{\mathfrak{p}\not\supseteq\mathfrak{b}_i} \left(1+\frac{1}{N(\mathfrak{p}^m)-1}\right) = \frac{1}{N(\mathfrak{b}_i^m)},$$

since the products combined give the Euler product for $\zeta_K(m)$. Alternatively, one can expand the products into sums and match terms with the series defining $\zeta_K(m)$. Now (26) and (27) imply that the multiple sum in (24) equals

$$\sum_{\mathfrak{b}_0}\sum_{\mathfrak{b}_1}\cdots\sum_{\mathfrak{b}_r} \frac{\mu(\mathfrak{b}_0)\mu(\mathfrak{b}_1)\cdots\mu(\mathfrak{b}_r)}{N(\operatorname{lcm}(\mathfrak{b}_0,\mathfrak{b}_1,\dots,\mathfrak{b}_r)^m)} E\begin{pmatrix} \mathfrak{b}_0^m & \mathfrak{b}_1^m & \dots & \mathfrak{b}_r^m \\ 0 & a_1 & \dots & a_r \end{pmatrix},$$

and by Lemma 2.6 we get the desired statement. □

3.1 Averages of the correlation functions

The following results will be useful in Section 4 when studying the spectral properties of the action $\mathcal{O}_K \curvearrowright (X,\Pi)$. Recall that the function g defined in (25) depends on m.

Lemma 3.4. *Let $\chi \in \Lambda'_{\mathfrak{d}^m}$. Then*

$$\lim_{x\to\infty} \frac{1}{\#B_x} \sum_{b\in B_x} \chi(b) c_2(b) = g^2(\mathfrak{d}).$$

Proof. Observe that

$$\lim_{x\to\infty} \frac{1}{\#B_x} \sum_{b\in B_x} \chi(b)\chi_1(b) = \begin{cases} 1 & \text{if } \chi_1 = \chi^{-1}; \\ 0 & \text{otherwise.} \end{cases} \tag{28}$$

By Lemma 3.3

$$\lim_{x\to\infty} \frac{1}{\#B_x} \sum_{b\in B_x} \chi(b) c_2(b) = \lim_{x\to\infty} \frac{1}{\#B_x} \sum_{b\in B_x} \chi(b) \sum_{\mathfrak{a}_0,\mathfrak{a}_1} g(\mathfrak{a}_0) g(\mathfrak{a}_1) \sum_{\substack{\chi_i \in \Lambda'_{\mathfrak{a}_i^m} \\ i=0,1 \\ \chi_0\chi_1 = \eta}} \chi_1(b) =$$

$$= \sum_{\mathfrak{a}_0,\mathfrak{a}_1} g(\mathfrak{a}_0) g(\mathfrak{a}_1) \sum_{\substack{\chi_i \in \Lambda'_{\mathfrak{a}_i^m} \\ i=0,1 \\ \chi_0\chi_1 = \eta}} \lim_{x\to\infty} \frac{1}{\#B_x} \sum_{b\in B_x} \chi(b)\chi_1(b) =$$

$$= g(\mathfrak{d}) \sum_{\mathfrak{a}_0} g(\mathfrak{a}_0) \sum_{\substack{\chi_0 \in \Lambda'_{\mathfrak{a}_0^m} \\ \chi_0\chi^{-1} = \eta}} 1 = g^2(\mathfrak{d}).$$

□

Lemma 3.5. *Let* $\chi_1 \in \Lambda'_{\mathfrak{d}_1^m}$, $\chi_2 \in \Lambda'_{\mathfrak{d}_2^m}$, *and* $\chi = \chi_1\chi_2 \in \Lambda'_{\mathfrak{d}^m}$. *Then*

$$\lim_{\substack{x \to \infty \\ y \to \infty}} \frac{1}{\#B_x \#B_y} \sum_{b_1 \in B_x} \sum_{b_2 \in B_y} \chi_1(b_1)\chi_2(b_2)c_3(b_1, b_2) = g(\mathfrak{d}_1)g(\mathfrak{d}_2)g(\mathfrak{d}). \tag{29}$$

Proof. Using (28) and Lemma 3.3 the LHS of (29) can be written as

$$\sum_{\mathfrak{a}_0,\mathfrak{a}_1,\mathfrak{a}_2} g(\mathfrak{a}_0)g(\mathfrak{a}_1)g(\mathfrak{a}_2) \sum_{\substack{\chi'_i \in \Lambda'_{\mathfrak{a}_i^m} \\ i=0,1,2 \\ \chi'_0\chi'_1\chi'_2 = \eta}} \lim_{\substack{x \to \infty \\ y \to \infty}} \frac{1}{\#B_x \#B_y} \sum_{b_1 \in B_x} \sum_{b_2 \in B_y} \chi_1(b_1)\chi_2(b_2)\chi'_1(b_1)\chi'_2(b_2) =$$

$$= \sum_{\mathfrak{a}_0} g(\mathfrak{a}_0)g(\mathfrak{d}_1)g(\mathfrak{d}_2) \sum_{\substack{\chi_0 \in \Lambda'_{\mathfrak{a}_0^m} \\ \chi'_0\chi_1^{-1}\chi_2^{-1} = \eta}} 1 = g(\mathfrak{d}_1)g(\mathfrak{d}_2)g(\mathfrak{d}).$$

$\square$

Notice that the proofs of Lemmata 3.4 and 3.5 are considerably simpler than the proofs of the analogous results in [3].

4 The Action $\mathcal{O}_K \curvearrowright (X, \Pi)$

In this Section we will prove the first part of our Main Theorem.

4.1 Ergodic and Pure Point Spectrum Actions

Let us recall the notion of ergodic and pure point spectrum action. Let G be a separable locally compact abelian[4] group and let S be a standard Borel G-space. Let μ be a σ-finite (left) G-invariant measure (that is $\mu(xE) = \mu(E)$ for all $x \in G$ and all Borel subset E of S). One says that the action $G \curvearrowright (S, \mu)$ is *ergodic* if whenever E is a Borel set in S with $\mu(E) \neq 0 \neq \mu(E^c)$, then for some $x \in G$ we have $\mu(E \vartriangle xE) > 0$. Let U be the unitary representation of G on $L^2(S, \mu)$ given by $(U_x f)(s) = f(xs)$. When U is a discrete direct sum of finite-dimensional irreducible representations one says that the action $G \curvearrowright (S, \mu)$ has *pure point spectrum*. This means that there exists an orthonormal basis $\{\Phi_j\}_{j=1}^{\infty}$ for $L^2(S, \mu)$ and a countable subgroup $\Gamma = \{\chi_j\}_{j=1}^{\infty}$ of $\hat{G}$ such that $U_x\Phi_j = \chi_j(x)\Phi_j$. The group Γ is referred to as the *spectrum* of the action.

One can construct ergodic actions with pure point spectrum as follows. Let K be a compact group, $H \subset K$ a closed subgroup, φ a continuous homomorphism of G onto a dense subgroup of K. Define the action $G \curvearrowright (K/H, \mu_H)$ by $x(k + H) = \varphi(x)k + H$, where μ_H is the unique measure on K/H such that $\mu_H(K/H) = 1$, G-invariant under left multiplication. One can check that the latter is ergodic (by transitivity of the action) and has pure point spectrum (the unitary representation U of G on $L^2(K/H, \mu_H)$ is given by $U_x = V_{\varphi(x)}$, where V is the unitary representation of K on $L^2(K/H, \mu_H)$; since V is a subrepresentation of the regular representation of K, it decomposes, by Peter-Weyl theorem, into a direct sum of finite-dimensional irreducible representations). Mackey [7] proved that, modulo removing null sets in S and K/H, every ergodic action with pure point spectrum can be realized as above. His work generalized the classical theory by von Neumann [16] and Halmos and von Neumann [5] where $G = \mathbf{Z}$. In all these cases, for actions with pure point spectrum, the isomorphism class is uniquely determined by its spectrum.

---

[4] The group G need not be abelian in order to define ergodic and pure point spectrum actions, see [7]. However, since in our case $G = \mathcal{O}_K \cong \mathbf{Z}^d$, it is not restrictive to make this assumption.

4.2 The Spectrum of the Action $\mathcal{O}_K \curvearrowright (X, \Pi)$

The goal of this section is to prove the following

Theorem 4.1. *The spectrum of the action* $\mathcal{O}_K \curvearrowright (X, \mathcal{B}, \Pi)$ *is* Λ.

For $a \in \mathcal{O}_K$, let U_a be the unitary operator on $\mathcal{H} = L^2(X, \Pi)$ given by

$$(U_a f)(x) = f(a \cdot x).$$

The proof of Theorem 4.1 requires to show that there exists an orthonormal basis $\{\theta_\chi\}_{\chi \in \Lambda}$ for $L^2(X, \Pi)$ such that $U_a \theta_\chi = \chi(a)\theta_\chi$. First we will show that Λ is contained in the spectrum of the action $\mathcal{O}_K \curvearrowright (X, \Pi)$. This is proven in the

Proposition 4.2. *Let* $\chi \in \Lambda$. *Then there exists a function* $\theta_\chi \in \mathcal{H}$ *such that*

$$(U_a \theta_\chi)(x) = \chi(a)\theta_\chi(x) \tag{30}$$

for Π-*almost every* $x \in X$.

Proof. Let $f_0 \in \mathcal{H}$, $f_0(x) = x(0)$, and for $a \in \mathcal{O}_K$ let $U_{a,\chi}$ be the unitary operator on $\mathcal{H}$ defined by

$$(U_{a,\chi} f)(x) = \chi(-a) f(a \cdot x).$$

Since $\mathcal{O}_K$ is amenable, von Neumann's mean ergodic theorem for $\mathcal{O}_K$-actions holds and implies that the limit

$$\lim_{R \to \infty} \frac{1}{\# B_R} \sum_{a \in B_R} U_{a,\chi} f_0$$

exists in $\mathcal{H}$. For Π-almost every $x \in X$, let us define

$$\begin{aligned} \lim_{R \to \infty} \frac{1}{\# B_R} \sum_{a \in B_R} U_{a,\chi} f_0(x) &= \lim_{R \to \infty} \frac{1}{\# B_R} \sum_{a \in B_R} \chi(-a) f_0(a \cdot x) = \\ &= \lim_{R \to \infty} \frac{1}{\# B_R} \sum_{a \in B_R} \chi(-a) x(a) =: \theta_\chi(x). \end{aligned} \tag{31}$$

Since θ_χ is $U_{a,\chi}$-invariant, i.e. $(U_{a,\lambda}\theta_\chi)(x) = \theta_\chi(x)$ for Π-almost every $x \in X$, we get that $\chi(-a)\theta_\chi(a \cdot x) = \theta_\chi(x)$, i.e. (30). $\square$

Notice that, since U_a is a unitary operator for every $a \in \mathcal{O}_K$, the eigenfunctions θ_χ are orthogonal to one another for different $\chi \in \Lambda$.

For $a \in \mathcal{O}_K$ let us denote by $x(a)$ the function $X \to \{0,1\}$ given by the projection of $x \in X$ onto its a-th coordinate. Introduce the subspace $H \subseteq \mathcal{H}$,

$$H = \overline{\left\{\sum_a z_a x(a)\right\}},$$

i.e. the closure of the set of all complex linear combinations of the $x(a)$'s. Notice that H is U_a-invariant for every $a \in \mathcal{O}_K$ and, by (31), all the eigenfunctions θ_χ belong to H. Let us write

$$x(a) = \sum_{\chi \in \Lambda} \langle x(a), \theta_\chi \rangle \theta_\chi.$$

Recall the function g from Lemma 3.3. We have the

Proposition 4.3. *Let $\chi \in \Lambda'_{\mathfrak{d}^m}$. Then for every $a \in \mathcal{O}_K$ we have*

$$\langle x(a), \theta_\chi \rangle = \zeta_K(m)\chi(a)g^2(\mathfrak{d}).$$

Proof. Observe that $\langle x(a), x(b) \rangle = \zeta_K(m)c_2(b-a)$. From (31) and Lemma 3.4 we get

$$\begin{aligned}\langle x(a), \theta_\chi \rangle &= \lim_{R\to\infty} \left\langle x(a), \frac{1}{|B_R|}\sum_{b\in B_R} \chi(-b)x(b) \right\rangle = \lim_{R\to\infty} \frac{1}{\#B_R}\sum_{b\in B_R} \chi(b)\langle x(a), x(b)\rangle = \\ &= \lim_{R\to\infty} \frac{1}{\#B_R}\sum_{b\in B_R} \chi(b)\zeta_K(m)c_2(b-a) = \zeta_K(m)\chi(a) \lim_{R\to\infty} \frac{1}{\#B_R}\sum_{b\in B_R} \chi(b)c_2(b) = \\ &= \zeta_K(m)\chi(a)g^2(\mathfrak{d}).\end{aligned}$$

$\square$

Since the inner products of θ_χ with all the functions $x(a)$, $a \in \mathcal{O}_K$, are positive, we immediately have the

Corollary 4.4. *All the eigenfunctions θ_χ are non-zero.*

An important step in the proof of Theorem 4.1 is given by the

Proposition 4.5. *The family of eigenfunctions $\{\theta_\chi\}_{\chi\in\Lambda}$ is a basis for H.*

Proof. By orthogonality, it is enough to show that the eigenfunctions span the space of all linear combinations of the $x(a)$'s. Let us show that H is isomorphic to $L^2(\widehat{\mathcal{O}_K}, \nu)$, where ν is the spectral measure.

The function $x \mapsto x(0)$ belongs to $L^2(X, \Pi)$ and for every $a \in \mathcal{O}_K$, we have $\langle U_a x(0), x(0)\rangle = c_2(a)$. Notice that

$$H = \overline{\text{span}\,\{U_a x(0) :\ a \in \mathcal{O}_K\}}.$$

By definition of ν, $\hat{\nu}(a) = c_2(a)$, that is

$$\int\limits_{\widehat{\mathcal{O}_K}} i(a)(\chi)\mathrm{d}\nu(\chi) = \langle U_a x(0), x(0)\rangle,$$

where $i : \mathcal{O}_K \to \widehat{\widehat{\mathcal{O}_K}}$ is the canonical isomorphism, $i(a)(\chi) = \chi(a)$. The map $L^2(X, \Pi) \to L^2(\widehat{\mathcal{O}_K}, \nu)$, $U_a x(0) \mapsto i(a)$ is an equivariant isometry (with respect to the $\mathcal{O}_K$-action). This extends to a unitary operator $W : H \to L^2(\widehat{\mathcal{O}_K}, \nu)$ and yields an isomorphism between the unitary representation $U|_H$ and V^ν, where $\mathcal{O}_K \ni a \mapsto (U|_H)_a := U_a|_H : H \to H$, $(U_a|_H f)(x) = f(a \cdot x)$ and $\mathcal{O}_K \ni a \mapsto V^\nu : L^2(\widehat{\mathcal{O}_K}, \nu) \to L^2(\widehat{\mathcal{O}_K}, \nu)$, $(V_a^\nu f)(\chi) = i(a)(\chi)f(\chi) = \chi(a)f(\chi)$. In particular, we have

$$H \cong L^2(\widehat{\mathcal{O}_K}, \nu) = \bigoplus_{\chi \in \Lambda} L^2(\widehat{\mathcal{O}_K}, \sigma_\chi \delta_\chi), \tag{32}$$

where

$$\sigma_\chi = \sum_{\mu^2(\mathfrak{d}) = 1 \,:\, \chi \in \Lambda_{\mathfrak{d}^m}} \sigma_{\mathfrak{d}}.$$

Since we have constructed a non-trivial eigenfunction θ_χ for each $\chi \in \Lambda$, (32) implies that the family of eigenfunctions $\{\theta_\chi\}_{\chi \in \Lambda}$ spans H. □

We want to normalize each eigenfunction to make the family $\{\theta_\chi\}_{\chi \in \Lambda}$ an orthonormal basis for H. For this purpose, we have the

Lemma 4.6. *Let $\chi \in \Lambda'_{\mathfrak{d}^m}$. Then*

$$\|\theta_\chi\| = \sqrt{\zeta_K(m)}|g(\mathfrak{d})|.$$

Proof. From Proposition 4.3 we get

$$\begin{aligned} \|\theta_\chi\|^2 &= \langle \theta_\chi, \theta_\chi\rangle = \left\langle \theta_\chi, \lim_{R\to\infty} \frac{1}{\#B_R} \sum_{a \in B_R} \chi(-a)x(a) \right\rangle = \lim_{R\to\infty} \frac{1}{\#B_R} \sum_{a \in B_R} \chi(a)\overline{\langle x(a), \theta_\chi\rangle} = \\ &= \lim_{R\to\infty} \frac{1}{\#B_R} \sum_{a \in B_R} |\chi(a)|^2 \zeta_K(m) g^2(\mathfrak{d}) = \zeta_K(m) g^2(\mathfrak{d}). \end{aligned}$$

□

Let us denote the the normalized eigenfunctions by $\tilde{\theta}_\chi = \theta_\chi/\|\theta_\chi\|$. If we write $x(a) = \sum_{\chi\in\Lambda}\langle x(a), \tilde{\theta}_\chi\rangle\tilde{\theta}_\chi$, then we can retrieve the fact that $\|x(a)\| = 1$ using Proposition 4.5 and Lemma 4.6:

$$\begin{aligned}
\|x(a)\| &= \sum_{\chi\in\Lambda}\left|\langle x(a), \tilde{\theta}_\chi\rangle\right|^2 = \sum_{\mu^2(\mathfrak{d})=1}\sum_{\chi\in\Lambda'_{\mathfrak{d}^m}}\zeta_K(m)g^2(\mathfrak{d}) = \\
&= \zeta_K(m)\sum_{\mu^2(\mathfrak{d})=1} g^2(\mathfrak{d})\#\Lambda'_{\mathfrak{d}^m} = \\
&= \frac{1}{\zeta_K(m)}\sum_{\mu^2(\mathfrak{d})=1}\prod_{\mathfrak{p}\supseteq\mathfrak{d}}\frac{1}{N(\mathfrak{p}^m)-1} = \frac{1}{\zeta_K(m)}\prod_{\mathfrak{p}}\left(1+\frac{1}{N(\mathfrak{p}^m)-1}\right) = \\
&= \frac{1}{\zeta_K(m)}\prod_{\mathfrak{p}}\left(1-\frac{1}{N(\mathfrak{p}^m)}\right)^{-1} = 1.
\end{aligned}$$

The same argument allows us to provide an approximation of the function $x(a)$ for $a \in \mathcal{O}_K$: let $D \geqslant 1$ and define

$$x_D(a) = \sum_{\substack{\mu^2(\mathfrak{d}) = 1 \\ N(\mathfrak{d}) \leqslant D}}\sum_{\chi\in\Lambda'_{\mathfrak{d}^m}}\left\langle x(a), \tilde{\theta}_\chi\right\rangle\tilde{\theta}_\chi.$$

We have the following estimate

$$\begin{aligned}
\|x(a) - x_D(a)\|^2 &= \sum_{\substack{\mu^2(\mathfrak{d}) = 1 \\ N(\mathfrak{d}) > D}}\sum_{\chi\in\Lambda'_{\mathfrak{d}^m}}\left|\left\langle x(a), \tilde{\theta}_\chi\right\rangle\right|^2 = \sum N(\mathfrak{d}) > D|g(\mathfrak{d})| = \\
&= O(D^{-1+\varepsilon}) \qquad (33)
\end{aligned}$$

for every $\varepsilon > 0$. Another important step in the proof of Theorem 4.1 is the

Proposition 4.7. *Let $\chi_1 \in \Lambda'_{\mathfrak{d}_1^m}$ and $\chi_2 \in \Lambda'_{\mathfrak{d}_2^m}$. Then*

$$\tilde{\theta}_{\chi_1}\tilde{\theta}_{\chi_2} = \varepsilon\tilde{\theta}_\chi,$$

where $\chi = \chi_1\chi_2 \in \Lambda'_{\mathfrak{d}^m}$ and $\varepsilon = \mu(\mathfrak{d}_1)\mu(\mathfrak{d}_2)\mu(\mathfrak{d})$.

Proof. It is enough to show that for every $a \in \mathcal{O}_K$ we have

$$\left\langle\tilde{\theta}_{\chi_1}\tilde{\theta}_{\chi_2}, x(a)\right\rangle = \varepsilon\left\langle\tilde{\theta}_\chi, x(a)\right\rangle.$$

Using the definition (31) we have

$$\theta_{\chi_1}\theta_{\chi_2} = \lim_{\substack{R_1\to\infty \\ R_2\to\infty}}\frac{1}{\#B_{R_1}\#B_{R_2}}\sum_{a_1\in B_{R_1}}\sum_{a_2\in B_{R_2}}\chi_1(-a_1)\chi_2(-a_2)x(a_1)x(a_2)$$

and thus

$$\begin{aligned}
&\langle \theta_{\chi_1}\theta_{\chi_2}, x(a)\rangle = \\
&= \lim_{\substack{R_1\to\infty\\ R_2\to\infty}} \frac{1}{\#B_{R_1}\#B_{R_2}} \sum_{a_1\in B_{R_1}}\sum_{a_2\in B_{R_2}} \chi_1(-a_1)\chi_2(-a_2)\,\langle x(a_1)x(a_2), x(a)\rangle = \\
&= \zeta_K(m) \lim_{\substack{R_1\to\infty\\ R_2\to\infty}} \frac{1}{\#B_{R_1}\#B_{R_2}} \sum_{a_1\in B_{R_1}}\sum_{a_2\in B_{R_2}} \chi_1(-a_1)\chi_2(-a_2)c_3(a_1-a, a_2-a) = \\
&= \zeta_K(m)(\chi_1\chi_2)(-a) \lim_{\substack{R_1\to\infty\\ R_2\to\infty}} \frac{1}{\#B_{R_1}\#B_{R_2}} \sum_{a_1\in B_{R_1}}\sum_{a_2\in B_{R_2}} \chi_1(-a_1)\chi_2(-a_2)c_3(a_1, a_2) = \\
&= \zeta_K(m)\chi(-a)g(\mathfrak{d}_1)g(\mathfrak{d}_2)g(\mathfrak{d})
\end{aligned}$$

by Lemma 3.5. On the other hand, by Proposition 4.3

$$\langle \theta_\chi, x(a)\rangle = \zeta_K(m)\chi(-a)g^2(\mathfrak{d}).$$

Therefore

$$\varepsilon = \left\langle \tilde\theta_{\chi_1}\tilde\theta_{\chi_2}, x(a)\right\rangle \left\langle \tilde\theta_\chi, x(a)\right\rangle^{-1} = \frac{g(\mathfrak{d}_1)g(\mathfrak{d}_2)g(\mathfrak{d})}{|g(\mathfrak{d}_1)||g(\mathfrak{d}_2)|}\frac{|g(\mathfrak{d})|}{g^2(\mathfrak{d})} = \mu(\mathfrak{d}_1)\mu(\mathfrak{d}_2)\mu(\mathfrak{d}).$$

□

We are now ready to prove the

Proposition 4.8. *$H = \mathcal{H}$.*

Proof. Let $\mathcal{B}$ denote the Borel σ-algebra on X. In the sense of Rokhlin [12], $(X, \mathcal{B}, \Pi)$ is a Lebesgue space. By Proposition 4.7, the space H is a sub-ring of the unitary ring $\mathcal{H} = L^2(X, \mathcal{B}, \Pi)$. Rokhlin's theorem [13] implies that $H = L^2(X, \mathcal{F}, \Pi|_{\mathcal{F}})$, where $\mathcal{F}$ is a σ-subalgebra of $\mathcal{B}$. We claim that, up to null sets, $\mathcal{F} = \mathcal{B}$. Let, by contradiction, assume that $\mathcal{F} \subsetneq \mathcal{B}$, i.e. there is a positive measure set in $\mathcal{B} \setminus \mathcal{F}$. The conditional expectation operator $\mathbf{E}(\cdot|\mathcal{F})$ is an orthogonal projection $\mathcal{H} \to H$, that is $\mathbf{E}(f|\mathcal{F}) = \mathrm{proj}_H(f)$ for every $f \in \mathcal{H}$. There exist a function $f \in \mathcal{H}$ and a constant $\alpha > 0$ such that $\|f - \mathbf{E}(f|\mathcal{F})\| \geqslant \alpha$. Let $\varepsilon > 0$, and let $f' = \sum_{i=1}^n \alpha_i \mathbf{1}_{A_i}$ be a simple function such that $\|f - f'\| \leqslant \frac{\varepsilon}{2}$, where A_i are cylinders (that is $A_i = \{x \in X : x(a_1^{(i)})x(a_2^{(i)})\cdots x(a_{r^{(i)}}^{(i)}) = 1\}$ for some $r^{(i)} \geqslant 1$ and $a_1^{(i)}, a_2^{(i)}, \ldots, a_{r^{(i)}}^{(i)} \in \mathcal{O}_K$). By (33) each function $x(a_j^{(i)})$ can be approximated arbitrarily well by a linear combination of the $\tilde\theta_\chi$'s and thus there exists a polynomial in the $\tilde\theta_\chi$'s, say f'', such that $\|f' - f''\| \leqslant \frac{\varepsilon}{2}$. By Proposition 4.7 the function f'' can be written as a *linear* combination of the $\tilde\theta_\chi$'s, and therefore, by Proposition 4.5, $f'' \in H$. This means that we are able to find $f \in H$ such that $\|f - f''\| \leqslant \varepsilon$ and, if ε is sufficiently small, this contradicts the fact that $\|f - \mathrm{proj}_H(f)\| \geqslant \alpha$. □

Propositions 4.5 and 4.8 immediately give the

Corollary 4.9. *The family of eigenfunctions* $\{\tilde{\theta}_\chi\}_{\chi \in \Lambda}$ *is an orthonormal basis for* $\mathcal{H}$.

This fact, together with Propositoin 4.2 and Corollary 4.4, yields Theorem 4.1, along with the

Theorem 4.10. *The action* $\mathcal{O}_K \curvearrowright (X, \Pi)$ *is ergodic.*

Theorems 4.1 and 4.10 constitute part (i) of our Main Theorem.

5 The action $\mathbf{Z}^d \curvearrowright (\mathbf{G}, \mathrm{Haar})$.

Recall the square-free solenoid $\mathbf{G}$ defined in (6) and the $\mathbf{Z}^d$ action given by (7). For $v \in \mathbf{Z}^d$ let $\mathbf{U}_v$ be the unitary operator on $\mathbf{H} = L^2(\mathbf{G}, \mathrm{Haar})$ given by

$$(\mathbf{U}_v f)(\mathbf{g}) = f(v \cdot \mathbf{g}).$$

We prove the

Proposition 5.1. *The spectrum of* $\mathbf{Z}^d \curvearrowright (\mathbf{G}, \mathrm{Haar})$ *is isomorphic to* Λ.

Proof. Let $\iota : \mathbf{Z}^d \to \mathcal{O}_K$ be the isomorphism defined in the Introduction. Let $\mathbf{g} \in \mathbf{G}$ and for every prime ideal $\mathfrak{p}$ let $g_{\mathfrak{p}^m} + \mathfrak{p}^m \in \mathcal{O}_K/\mathfrak{p}^m$ be its projection onto the $\mathfrak{p}^m$-th coordinate. Let $\chi \in \Lambda'_{\mathfrak{p}^m}$. Notice that if $a \equiv a' \mod \mathfrak{p}^m$, then $\chi(a) = \chi(a')$; in other words, χ is well defined on $\mathcal{O}_K/\mathfrak{p}^m$. Let $\xi_\chi(\mathbf{g}) = \chi(g_{\mathfrak{p}^m} + \mathfrak{p}^m)$. It is clear that $(\mathbf{U}_v \xi_\chi)(\mathbf{g}) = \chi(\iota(v))\xi_\chi(\mathbf{g})$, i.e. ξ is an eigenfunction with eigenvalue $\chi(\iota(v))$. If $\chi \in \Lambda'_{\mathfrak{d}^2}$ and $\mathfrak{d} = \mathfrak{p}_1 \cdots \mathfrak{p}_s$ (distinct prime ideals), then $\chi = \chi_1 \cdots \chi_s$, where $\chi_i \in \Lambda'_{\mathfrak{p}_i^m}$; in this case the function $\xi_\chi(\mathbf{g}) = \chi_1(g_{\mathfrak{p}_1^m} + \mathfrak{p}_1^m) \cdots \chi_s(g_{\mathfrak{p}_s^m} + \mathfrak{p}_s^m)$ is an eigenfunction for $\mathbf{U}_v$ with eigenvalue $\chi(\iota(v))$. Since characters are orthonormal with respect to the Haar measure on $\mathbf{G}$, we have that the discrete group $\{\chi \circ \iota\}_{\chi \in \Lambda} \subseteq \widehat{\mathbf{Z}^d} = \mathbf{T}^d$ is the spectrum of the action $\mathbf{Z}^d \curvearrowright (\mathbf{G}, \mathrm{Haar})$ and is clearly isomorphic to Λ. $\square$

The above proposition and Mackey's theorem [7] yield the

Corollary 5.2. *The two actions* $\mathcal{O}_K \curvearrowright (X, \Pi)$ *and* $\mathbf{Z}^d \curvearrowright (\mathbf{G}, \mathrm{Haar})$ *are isomorphic.*

The above corollary constitutes part (ii) of our Main Theorem. Corollary 1.4 from the Introduction follows immediately from our Main Theorem: see, e.g. [2] to get absence of weak mixing and [17] to get zero measure-theoretical entropy.

6 Example: Square-Free Gaussian Integers

Let $K = \mathbf{Q}(i)$ and $m = 2$. Then $\mathcal{O}_K = \mathbf{Z}[i]$ is given by Gaussian integers. Since $\mathbf{Z}[i]$ is a principal ideal domain, every ideal is of the form $(a + ib)$ for some $a, b \in \mathbf{Z}$. The norm is given by $N((a + ib)) = a^2 + b^2$. Units in $\mathcal{O}_K$ are $\pm 1, \pm i$. Prime ideals $\mathfrak{p}$ are of the form

- (p), where $p \in \mathbf{Z}$ is a usual prime and $p \equiv 3 \bmod 4$,
- $(a + ib)$, such that $a^2 + b^2 \in \mathbf{Z}$ is a usual prime (necessarily equal to 2 or congruent to $1 \bmod 4$).

The prime ideals $\mathfrak{p}$ with norm $N(\mathfrak{p}) \leqslant 13$ are shown, together with their squares, in Figure 1. Square-free Gaussian integers are shown Figure 2.



Figure 1: Prime ideals (red) in $\mathbf{Z}[i]$ with norm at most 13, and their squares (blue). We have $N((1 + i)) = 2$, $N((1 \pm 2i)) = 5$, $N((3)) = 9$, $N((2 \pm 3i)) = 13$.



Figure 2: Square-free Gaussian integers. The square grid is identified with $\mathbf{Z}[i] \cong \mathbf{Z}^2$ and each square is colored black (resp. white) if it corresponds to a square-free (resp. not square-free) Gaussian integer. On the left: $\mathcal{F}_2 \cap \{a+ib\}$ where $-50 \leqslant a, b \leqslant 50$. Notice the dihedral D_4 symmetry. On the right, $\mathcal{F}_2 \cap \{a+ib\}$ where $10^{12} \leqslant a \leqslant 10^{12}+100$ and $10^{15} \leqslant b \leqslant 10^{15}+100$.

In this case we can write the Dedekind zeta function in terms of primes in $\mathbf{Z}$. For $\Re s > 1$ we have

$$\begin{aligned}\zeta_{\mathbf{Q}(i)}(s) &= \prod_{\mathfrak{p}} \bigl(1 - N^{-s}(\mathfrak{p})\bigr) = \\ &= (1-2^{-s})^{-1} \prod_{p \equiv 1 \bmod 4} (1-p^{-s})^{-2} \prod_{p \equiv 3 \bmod 4} (1-p^{-2s})^{-1} = \zeta(s)\beta(s),\end{aligned}$$

where ζ denotes the Riemann zeta function and β the Dirichlet beta function,

$$\beta(s) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s}.$$

The density of square-free Gaussian integers is $1/\zeta_{\mathbf{Q}(i)}(2) = \frac{6}{\pi^2 G} \approx 0.6637$, where $G = \beta(2)$ is the Catalan constant.

Let us look at correlation functions. For example, $c_4(1, i, 1+i) = 0$ because for every $a \in \mathbf{Z}[i]$ at least one of the four Gaussian integers a, $a+1$, $a+i$, $a+1+i$, is not square-free, since it is divisible by $2 = (1+i)^2$. This is reflected by the formula (3), where $D(\mathfrak{p}^2 \mid 0, 1, i, 1+i) = 4$ for all prime ideals $\mathfrak{p}$ and there is a prime ideal, $\mathfrak{p}_1 = (1+i)$, for which $N(\mathfrak{p}_1^2) = 4$. On the other hand, $c_5(1,i,-1,-i) = \left(1-\frac{3}{4}\right)\prod_{\mathfrak{p}\neq\mathfrak{p}_1}\left(1-\frac{5}{N(\mathfrak{p}^2)}\right) > 0$, since $\pi_{\mathfrak{p}_1^2}(\{0,1,i,-1,-i\}) = \{0+\mathfrak{p}_1^2, 1+\mathfrak{p}_1^2, i+\mathfrak{p}_1^2\}$, and for every prime ideal $\mathfrak{p}\neq\mathfrak{p}_1$ we have $D(\mathfrak{p}^2 \mid 0,1,i,-1,-i) = 5$ and $N(\mathfrak{p}) \geqslant 5$. More precisely

$$c_5(1,i,-1,-i) = \frac{1}{4}\prod_{p\equiv 1 \mod 4}\left(1-\frac{5}{p^2}\right)^2 \prod_{p\equiv 3 \mod 4}\left(1-\frac{5}{p^4}\right) \approx 0.1303.$$



Figure 3: On the left: the fundamental domain for the ideal $(8-6i) = (1+i)^2(1-2i)^2$. It contains $N(8-6i) = 2^2 5^2 = 100$ integers. On the right, $\Lambda_{(8-6i)}$.

Let us describe the spectrum Λ of the $\mathbf{Z}[i]$-action on (X,Π). By choosing the isomorphism $\iota : \mathbf{Z}^2 \to \mathbf{Z}[i]$, $(a,b)\mapsto a+ib$, we can identify Λ with a subset of $\mathbf{T}^2$. For every square-free ideal $\mathfrak{d}\subseteq\mathbf{Z}^2$, let us consider a fundamental domain $\mathfrak{F}_{\mathfrak{d}^2}$ for $\mathbf{Z}^2/\mathfrak{d}^2$, say the square with sides $w=(w_1,w_2)$, and $w'=(-w_2,w_1)$, where $w_1>0$, $w_2\geqslant 0$, and $w_1^2+w_2^2 = N(\mathfrak{d})$. In this way $\#\mathfrak{F}_{\mathfrak{d}^2} = N(\mathfrak{d}^2)$. One can check that

$$\Lambda_{\mathfrak{d}^2} = \frac{1}{N(\mathfrak{d}^2)}\begin{pmatrix} -w_2 & w_1 \\ w_1 & w_2 \end{pmatrix}\mathfrak{F}_{\mathfrak{d}^2} \subseteq \left\{\left(\frac{t_1}{N(\mathfrak{d}^2)},\frac{t_2}{N(\mathfrak{d}^2)}\right)\in\mathbf{T}^2,\ 0\leqslant t_1,t_2 < N(\mathfrak{d}^2)\right\}.$$

See Figure 3 for an example.

The spectrum Λ is the subgroup of $\mathbf{T}^2$ obtained as union of the $\Lambda_{\mathfrak{d}^2}$'s as above, and is shown in Figure 4.



Figure 4: Successive approximations $\bigcup_{N(\mathfrak{d}^2)\leqslant D}\Lambda_{\mathfrak{d}^2}$ of the spectrum Λ, with $D = 25, 100, 625, 2809$. The size of points in $\Lambda_{\mathfrak{d}^2}$ decreases as $N(\mathfrak{d}^2)$ increases.

References

[1] M. Baake, R.V. Moody, and P.A.B. Pleasants. Diffraction from visible lattice points and kth power free integers. *Discrete Math.*, 221(1-3):3–42, 2000. Selected papers in honor of Ludwig Danzer.

[2] V. Bergelson and A. Gorodnik. Weakly mixing group actions: a brief survey and an example. In *Modern dynamical systems and applications*, pages 3–25. Cambridge Univ. Press, Cambridge, 2004.

[3] F. Cellarosi and Ya.G Sinai. Ergodic properties of square-free numbers. *to appear in* Journal of the European Mathematical Society, http://arxiv.org/abs/1112.4691.

[4] R.R. Hall. The distribution of squarefree numbers. *J. Reine Angew. Math.*, 394:107–117, 1989.

[5] P.R. Halmos and J. von Neumann. Operator methods in classical mechanics. II. *Ann. of Math. (2)*, 43:332–350, 1942.

[6] D.R. Heath-Brown. The square sieve and consecutive square-free numbers. *Math. Ann.*, 266(3):251–259, 1984.

[7] G. W. Mackey. Ergodic transformation groups with a pure point spectrum. *Illinois J. Math.*, 8:593–600, 1964.

[8] L. Mirsky. Arithmetical pattern problems relating to divisibility by rth powers. *Proc. London Math. Soc. (2)*, 50:497–508, 1949.

[9] W. Narkiewicz. *Elementary and analytic theory of algebraic numbers.* Springer Monographs in Mathematics. Springer-Verlag, Berlin, third edition, 2004.

[10] J. Neukirch. *Algebraic number theory*, volume 322 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.

[11] P.A.B. Pleasants and C. Huck. Entropy and diffraction of the k-free points in the n-dimensional lattice. *to appear.*

[12] V.A. Rokhlin. On the problem of the classification of automorphisms of Lebesgue spaces. *Doklady Akad. Nauk SSSR (N. S.)*, 58:189–191, 1947.

[13] V.A. Rokhlin. Unitary rings. *Doklady Akad. Nauk SSSR (N.S.)*, 59:643–646, 1948.

[14] P. Sarnak. Three lectures on the Möbius function randomness and dynamics (Lecture 1). `http://publications.ias.edu/sites/default/files/MobiusFunctionsLectures(2).pdf`.

[15] K.M. Tsang. The distribution of r-tuples of squarefree numbers. *Mathematika*, 32(2):265–275 (1986), 1985.

[16] J. von Neumann. Zur Operatorenmethode in der klassischen Mechanik. *Ann. of Math. (2)*, 33(3):587–642, 1932.

[17] R. J. Zimmer. Ergodic actions with generalized discrete spectrum. *Illinois J. Math.*, 20(4):555–588, 1976.